


SOD ANAMIX[®] EARLY YEARS

A methionine- and cystine-free powdered infant formula with iron containing a balanced blend of essential and non-essential amino acids, carbohydrate (including prebiotics), fat (including DHA and ARA), vitamins, minerals and trace elements.

Must Be Used Under Medical Supervision



Indications: Sulfite Oxidase Deficiency (SOD)

Suitable for infants and young children.


Ordering/Reimbursement Information for SOD Anamix Early Years

Product Code	Reimbursement Code	HCPCS Code*	Product Packaging	Calories Per Can
90212 (Special Order)	49735-0102-12	B4162	3 x 400 g (14.1 oz)	1892

* S9435 = For individuals who submit their bills to insurance on their own.

Mixing Instructions:

- To make a standard dilution (21 kcal/fl oz), add 1 unpacked level scoop (5 g) of powder to each fluid ounce of water. Only use the scoop provided. For accuracy formula should be weighed on a gram scale.
- SOD Anamix Early Years may be mixed with warm or cool water that has been boiled. Do not boil formula or use a microwave to prepare or warm formula.



To make formula:

1. Wash hands thoroughly and clean the preparation area.
2. Pour the required amount of cooled, boiled water into the sterilized* feeding bottle.
3. Add the prescribed number of leveled scoops of formula to the water in the sterilized feeding bottle.
4. Place cap on the bottle and shake until the powder dissolves. Shake immediately before use.

*Ask your baby's health care team if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

Important Preparation Information:

Freshly prepared formula is best. Formula prepared in advance must be cooled rapidly, stored in the refrigerator and used within 24 hours of preparation. Formula remaining in the bottle after one hour of feeding should be discarded. Powdered infant formula is not sterile.

Storage:

Store unopened cans in a cool, dry place. Avoid extreme temperatures. Once opened, use contents within 2 weeks. Replace lid firmly after use.

Important Notice:

Must only be under medical supervision. For oral and tube feeding. Not for parenteral use. Not suitable as a sole source of nutrition. Only suitable for infants and young children with proven Sulfite Oxidase Deficiency (SOD) or other conditions requiring restriction of methionine and cystine.

Amount of Water	Amount of SOD Anamix Early Years Powder to Add	Approximate Protein Equivalent	Approximate Final Volume
5 fl oz	25 g (approx. 5 scoops)	3.4 g	5.6 fl oz
8 fl oz	40 g (approx. 8 scoops)	5.4 g	9 fl oz
12 fl oz	60 g (approx. 12 scoops)	8.1 g	13.5 fl oz